

NEAR-INFRARED PHOTOMETRY AND SPECTROSCOPY OF L AND T DWARFS: THE EFFECTS OF CLOUDS, GRAVITY, AND EFFECTIVE TEMPERATURE

D. A. Golimowski¹, S. K. Leggett², M. S. Marley³, X. Fan⁴, T. R. Geballe⁵, and G. R. Knapp⁶

¹Department of Physics and Astronomy, Johns Hopkins University, 3400 N. Charles St., Baltimore, MD 21218

²United Kingdom Infrared Telescope, Joint Astronomy Centre, 660 N. A'ohoku Place, Hilo, HI 96720

³NASA-Ames Research Center, Mail Stop 245-3, Moffett Field, CA 94035

⁴Steward Observatory, University of Arizona, Tucson, AZ 85721

⁵Gemini Observatory, 670 N. A'ohoku Place, Hilo, HI 96720

⁶Princeton University Observatory, Princeton, NJ 08544

ABSTRACT

We present new $JHKLM'$ photometry on the MKO system for a large sample of L and T dwarfs identified from SDSS and 2MASS and classified according to the scheme of Geballe et al. (2002). We have compiled a sample of 105 L and T dwarfs that are uniformly classified and measured on a single photometric system. The scattered JHK spectral indices and colors of L dwarfs are likely caused by variations in the altitudes, distributions, and thicknesses of condensate clouds. Scatter in the $H-K$ colors of late T dwarfs probably reflects the sensitivity of the K -band flux to pressure induced H_2 opacity, which itself is sensitive to surface gravity. The M' luminosities of late-T dwarfs are 1.5–2.5 times fainter than predicted under conditions of chemical equilibrium. We have computed L_{bol} and T_{eff} for 42 L and T dwarfs whose trigonometric parallaxes have been measured. We find that $T_{\text{eff}} \approx 1450$ K for types L7–T4, which supports recent models that attribute the changing JHK -band luminosities and spectral features across the L–T transition to rapid changes in the condensate clouds over a narrow range of T_{eff} . We compute $T_{\text{eff}} = 600\text{--}750$ K for 2MASS 0415-0935 (T9), which supplants Gl 570D as the coolest known brown dwarf.

Key words: infrared: stars – stars: fundamental parameters – stars: late-type – stars: low-mass, brown dwarfs




Figure 1. Near-IR colors of L and T dwarfs. Typical error bars are shown. Encircled points represent the unusually blue late-L dwarfs SDSS J0805+4812, SDSS J0931+0327, SDSS J1104+5548, and SDSS J1331-0116.

1. PHOTOMETRIC DATA

We have compiled complete or partial $1\text{--}5\ \mu\text{m}$ photometry for 105 spectroscopically-confirmed L and T dwarfs using United Kingdom Infrared Telescope (UKIRT) imaging cameras and UKIRT Z and Mauna Kea Observatory (MKO) $JHKLM'$ filters. Diagrams of color versus near-infrared spectral type (Geballe et al. 2002) are shown in Figure 1 for various combinations of bandpasses.

The Z through K colors of L dwarfs are scattered because their $1\text{--}2.5\ \mu\text{m}$ “photospheres” coincide with clouds of Fe and silicate condensates with varying optical depths (Ackerman & Marley 2001, Marley et al. 2002, Tsuji 2002). Thus, near-IR colors are not good indicators of L subtype. On the other hand, $J-H$ is a good indicator of T

subtype presumably because, for $T_{\text{eff}} < 1400$ K, the condensate clouds have settled below the near-infrared photosphere. Conversely, $H-K$ is broadly scattered among T dwarfs because the K flux is affected by pressure-induced H_2 absorption, which is very sensitive to surface gravity (Burgasser et al. 2002, and references therein). The $Z-J$ colors of T dwarfs are also scattered, possibly because of variable process(es) responsible for removing condensates at the L–T transition, or because the Z flux is affected by the gravity- and metallicity-sensitive wings of the Na-D and K resonance lines.

The $K-L'$ and $L'-M'$ colors of L6–T4 dwarfs are nearly constant. The constancy of $K-L'$ over this range of types is explained by balanced CH_4 absorption in the K and L'

bands and the redistribution of continuum flux as the condensate clouds settle. On the other hand, $L'-M'$ is largely unaffected by the settling clouds. Its constancy for types L6–T4 reflects the constancy of T_{eff} throughout the L–T transition. (See §3.)

2. SPECTROSCOPIC DATA

We have obtained JHK spectra of 23 new L dwarfs and 14 new T0–T7 dwarfs identified from the Sloan Digital Sky Survey (SDSS) photometric database using UKIRT’s CGS4 and UIST spectrographs. We have also obtained new spectra of 11 L and T dwarfs previously identified from the Two-Micron All Sky Survey (2MASS), including the coolest known T dwarf, 2MASS J0415–0935. We have classified these dwarfs according to the scheme of Geballe et al. (2002). Our augmented sample shows that some spectral indices of this scheme are not internally consistent for L dwarfs because of their apparent sensitivity to cloud optical depth. Some adjustment of these indices is warranted.

Figure 2 shows the H and K spectra of representative T6, T7, T8, and T9 dwarfs. These spectra show a steady increase in the depths of the H_2O and CH_4 bands with increasing spectral type. The relatively small spectral differences between the dwarfs designated T8 and T9 may be more indicative of a half-subtype increment, rather than a whole-subtype increment. Regardless, this sequence suggests that there is room for one more subtype, which would have essentially zero flux at 1.45, 1.7, and 2.25 μm .

The effects of varying cloud optical depth and gravity are manifested not only in the broadband colors of L and T dwarfs (§1), but in the equivalent widths of nar-




Figure 2. Normalized H and K spectra ($R \approx 600$) for (top to bottom) SDSS J1624+0029 (T6), SDSS J1758+4633 (T7), Gl 570D (T8), and 2MASS J0415–0935 (T9).




Figure 3. Normalized J spectra ($R \approx 600$) for 3 of the 4 unusually blue late-L dwarfs, bracketed by more typical L5.5 and L9 dwarfs. Strong FeH, K I, and H_2O absorption features are identified.

rower atomic and molecular absorption lines. Figure 3 compares the J spectra of three unusually blue late-L dwarfs (marked with circles in Figure 1) with those of more typical L dwarfs. The enhanced strengths of the FeH and K I features in the former group mimic those of earlier L dwarfs, which suggests that either the condensate clouds of these late-L dwarfs are unusually thin, or these dwarfs have low metallicity.

Figure 4 shows J spectra of several T6 and T8 dwarfs that exhibit progressively stronger absorption by K I at 1.243 and 1.254 μm . Recent investigations of spectroscopic gravity indicators in T dwarfs indicate that this doublet strengthens with decreasing gravity (Martín & Zapaterio Osorio 2002). This trend is supported by the increasing $H-K$ colors of these dwarfs (see Figure 7), which reflect progressively less




Figure 4. Normalized J spectra ($R \approx 600$) for T6 (left) and T8 (right) dwarfs. The K I absorption strengths and the $H-K$ colors of these dwarfs increase from top to bottom. Both trends indicate decreasing surface gravity.

pressure-induced H_2 absorption of K -band flux as gravity decreases (Burgasser et al. 2002).

3. BOLOMETRIC CORRECTIONS AND T_{eff}

We used our $ZJHK$ spectra, our $ZJHKL'$ photometry, and recently published parallaxes (Dahn et al. 2002, Tinney, Vrba et al. 2004) to compute BC_K and T_{eff} for late-M, L, and T dwarfs. In doing so, we assumed a Rayleigh-Jeans flux distribution longward of L' appropriately corrected for CH_4 and CO absorption between 3 and 5 μm . We used available M' photometry to confirm the accuracy of these assumptions. To compute T_{eff} , we adopted the ranges of brown-dwarf radii predicted by the evolutionary models of Burrows et al. (1997), Baraffe et al. (1998), and Chabrier et al. (2000) for ages 0.1–10 Gyr. (Narrower ranges were used for those dwarfs whose ages have been spectroscopically constrained.) Nominal values of T_{eff} for each dwarf were computed assuming a mean age of 3 Gyr for the solar neighborhood (Dahn et al. 2002).

Figure 5 shows the behavior of BC_K and T_{eff} as a function of spectral type. BC_K rises from M5 to L5 due to Wien shift and enhanced K luminosity as clouds settle below the K photosphere. It declines after L5 because of increasing CH_4 absorption at 2.3 μm . The scatter among late-T dwarfs shows the sensitivity of H_2 absorption to gravity variations. T_{eff} declines steeply and monotonically for types M6–L7 and T4–T9, but is nearly constant (~ 1450 K) over the L–T transition. This constancy indicates that the condensate clouds settle, thin, or disintegrate over a small range of T_{eff} . Thus, the presently defined L and T spectral classes are not proxies for a uniformly decreasing T_{eff} scale. We compute $T_{\text{eff}} = 600$ –




Figure 5. Diagrams of (a) BC_K and (b) T_{eff} versus spectral type for ultracool dwarfs. The plotted values of T_{eff} correspond to an age of 3 Gyr, and the error bars reflect the range of T_{eff} for ages 0.1–10 Gyr. The curves are polynomial fits to the data whose coefficients are reported by Golimowski et al. (2004).

750 K for the T9 dwarf 2MASS 0415-0935, which supplants Gl 570D as the coolest known brown dwarf.

4. COMPARISON WITH PRECIPITATING CLOUD MODELS

We compare our absolute photometry with the magnitudes and colors of L and T dwarfs predicted by the precipitating cloud models of Ackerman & Marley (2001) and Marley et al. (2002). The models are parametrized by f_{sed} , which describes the sedimentation efficiency relative to the upward transport of condensates by convection. Figure 6 shows M_J as functions of spectral type and $J-K$ color. A polynomial fit to the data in the left panel emphasizes the brightening J luminosity between types T0 and T4. This “early-T” hump (Tinney, Burgasser, & Kirkpatrick 2003) is probably caused by the thinning, settling, and/or break-up of the cloud deck, which allows J flux to emerge from deeper layers of the atmosphere.

Overplotted in the right panel of Figure 6 are model sequences for $f_{\text{sed}} = 3, 5$, and ∞ (no clouds) and different gravities ($\log g$). The $f_{\text{sed}} = 3$ models match the L dwarf data well, but do not turn fast enough to match the T dwarf data, which are better matched by the cloudless models. The dotted lines connect the $T_{\text{eff}} = 1300$ K points (triangles) on each model sequence. These lines bound the L–T transition, which suggests that holes in the cloud deck develop rapidly over a narrow range of T_{eff} . This notion is supported by the nearly constant $T_{\text{eff}} \approx 1450$ K measured bolometrically for types L7–T4 (Figure 5).

Figure 7 is a $J-H$ versus $H-K$ diagram for T dwarfs overlaid with model sequences for various combinations




Figure 6. Plots of M_J versus spectral type and $J-K$. Error bars are shown where the uncertainties in the distance modulus and spectral type are ≥ 0.1 mag and > 1 subclass, respectively. Known binaries are encircled. Model sequences are shown in the right panel for different values of f_{sed} and $\log g$.




Figure 7. Plots of $J-H$ vs. $H-K$ for T dwarfs. Color sequences are shown for $f_{\text{sed}} = 5$, $\log g = 5.0$ (dashed curve) and cloudless, $\log g = 5.5, 5.0, 4.5$, and 4.0 (solid curves, left to right) models. T_{eff} is marked in units of 100 K; dotted lines represent constant T_{eff} . A typical error bar is shown.

of f_{sed} and gravity. Because the radii of brown dwarfs older than 200 Myr vary by $< 30\%$ (Marley et al. 1996; Burrows et al. 2001), gravity is tightly correlated with mass. Thus the solid curves in Figure 7 are effectively evolutionary tracks of brown dwarfs with masses of (left to right) 75, 35, 15, and 10 M_{Jup} . Consequently, the T5.5 dwarf SDSS J1110+0116 may be a 10–15 M_{Jup} brown dwarf. As stated in §1, the K fluxes of T dwarfs are highly sensitive to gravity-dependent, pressure-induced H_2 absorption. Thus, $H-K$ is a good indicator of mass, such that bluer $H-K$ reflects higher mass for a given T subtype.

5. EFFECTS OF NONEQUILIBRIUM CHEMISTRY

The model atmospheres of Marley et al. (2002) assume all species are in thermochemical equilibrium. However, the M -band spectrum of the archetypal T dwarf GJ 229B shows absorption at $4.7 \mu\text{m}$ by overabundant CO dredged upward from hotter regions by convective mixing (Noll, Geballe & Marley 1997; and Oppenheimer et al. 1998). The same phenomenon should occur in other late-T dwarfs.

Figure 8 compares the cloudy and cloud-free models of Marley et al. (2002) with our measurements of M_K , $M_{L'}$, and $M_{M'}$ versus T_{eff} for L and T dwarfs. The $\log g = 5.0$ – 5.5 cloudy models match M_K , $M_{L'}$, and $M_{M'}$ well for L and early-T dwarfs. The $\log g = 4.5$ – 5.0 cloud-free models generally match M_K and $M_{L'}$ for late-T dwarfs, but they overestimate the M' luminosities of these dwarfs by factors of 1.5 to 2.5. We attribute these M' flux deficits to nonequilibrium abundances of CO, as shown by the vertical-mixing models of Saumon et al. (2003). These over-predicted M' fluxes affect the expected sensitivities of $5 \mu\text{m}$ searches for even cooler, “infra-T” dwarfs, as are planned with the *Spitzer Space Telescope*.




Figure 8. Plots of M_K , $M_{L'}$, and $M_{M'}$ vs. T_{eff} for L and T dwarfs, overlayed with predicted sequences from equilibrium models with $f_{\text{sed}} = 3$ (green), $f_{\text{sed}} = 5$ (red), and cloud-free (blue) atmospheres and surface gravities of $\log g = 4.5$ (short dash), 5.0 (solid), and 5.5 (long dash).

ACKNOWLEDGEMENTS

We thank Didier Saumon for computing the model magnitudes shown in Figures 6–8. This work is more completely described by Knapp et al. (2004) and Golimowski et al. (2004). Tables of data and spectra are available at <http://www.jach.hawaii.edu/~skl/LTdata.html>.

REFERENCES

- Ackerman, A. S., & Marley, M. S. 2001, ApJ, 556, 872
- Baraffe, I., et al. 1998, A&A, 337, 403
- Burgasser, A. J., et al. 2002, ApJ, 564, 421
- Burrows, A., Marley, M., et al. 1997, ApJ, 491, 856
- Burrows, A., et al. 2001, Rev. Mod. Phys., 73, 719
- Chabrier, G., et al. 2000, ApJ, 542, 464
- Dahn, C. C., et al. 2002, AJ, 124, 1170
- Geballe, T. R., et al. 2002, ApJ, 564, 466
- Golimowski, D. A., et al. 2004, AJ, 127, 3516
- Knapp, G. R., et al. 2004, AJ, 127, 3553
- Marley, M. S., et al. 1996, Science, 272, 1919
- Marley, M. S., et al. 2002, ApJ, 568, 335

- Martín, E. L., & Zapaterio Osorio, M. R. 2003, ApJ, 593, L113
- Noll, K. S., Geballe, T. R., & Marley, M. S. 1997, ApJ, 489, L87
- Oppenheimer, B. R., et al. 1998, ApJ, 502, 932
- Saumon, D., et al. 2003, in IAU Symp. 211, Brown Dwarfs, ed.
E. Martín (San Francisco: ASP), 345
- Tinney, C., Burgasser, A., & Kirkpatrick, J. 2003, AJ, 126, 975
- Tsuji, T. 2002, ApJ, 575, 264
- Vrba, F. J., et al. 2004, AJ, 127, 2948